INTERNATIONAL CANINE SEMEN BANK - OREGON MAIN OFFICE | P.O. Box 651|Sandy, Oregon 97055 Phone: 503-663-7031 | FAX: (503) 676-8025

Email: ik9sb@aol.com | Website: www.ik9sb.com

ICSB Canine Semen Volume Extension Media

The purpose of the ICSB Canine Semen Volume Extension Media is threefold. The media can:

- Increase the volume of low-volume ejaculates to provide a greater likelihood of conception.
- Increase the general quality of the ejaculate by improving sperm motility and progressive movement, thereby giving the sperm cells a safe boost of energy and nutrients.
- Provide an excellent medium for extending the volume of the dog's ejaculate where the semen is to be divided for breeding multiple bitches at the same time.

In order to use the ICSB Canine Semen Volume Extension Media, it is necessary to keep the media frozen until just prior to use. When needed, thaw the media at room to body temperature (70-100°F) and then add the desired amount of the media to the collected semen sample. Gently and thoroughly mix the sample with the media for best results.

Please contact ICSB if you have any questions or concerns!



INTERNATIONAL CANINE SEMEN BANK - OREGON MAIN OFFICE | P.O. Box 651|Sandy, Oregon 97055 Phone: 503-663-7031 | FAX: (503) 676-8025 Email: ik9sb@aol.com | Website: www.ik9sb.com

ICSB Canine Semen Volume Extension Media

The purpose of the ICSB Canine Semen Volume Extension Media is threefold. The media can:

- Increase the volume of low-volume ejaculates to provide a greater likelihood of conception.
- Increase the general quality of the ejaculate by improving sperm motility and progressive movement, thereby giving the sperm cells a safe boost of energy and nutrients.
- Provide an excellent medium for extending the volume of the dog's ejaculate where the semen is to be divided for breeding multiple bitches at the same time.

In order to use the ICSB Canine Semen Volume Extension Media, it is necessary to keep the media frozen until just prior to use. When needed, thaw the media at room to body temperature (70-100°F) and then add the desired amount of the media to the collected semen sample. Gently and thoroughly mix the sample with the media for best results.

Please contact ICSB if you have any questions or concerns!